

Research Article
Open Access

Growth Performance, Carcass Characteristics, Immune Response and Meat Quality of Broiler Chickens Fed *Muntingia Calabura* Leaf Extract

Alagbe Olujimi John

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India

ABSTRACT

This experiment was carried out to determine the effect of feeding *Muntingia calabura* leaf extract on the growth performance, carcass characteristics, immune response and meat quality of broiler chickens. 500 1-day old broiler chicks (Cobb 500) were randomly distributed into five treatments of five replicates consisting of ten birds each in a completely randomized design model. Birds in treatment one (control) was fed basal diet alone which is adequate in all nutrients according to the recommendation of broilers while treatment two, three, four and five were fed same diet with *Muntingia calabura* leaf extract at 0.3 mL, 0.6 mL, 0.9 mL and 1.2 mL per liter of water correspondingly. The duration of the experiment was 56 days and birds had unrestricted access to feed and fresh water. Body weight gain was higher in birds fed treatment four (2555.3 g) and five (2646.1 g), intermediate in treatment two (2346.6 g) and three (2365.1 g) and lower in treatment one (1948.8 g) ($P < 0.05$). Total feed intake and dressing percentage was higher among birds fed *Muntingia calabura* leaf extract relative to the control ($P < 0.05$). Conversely, mortality of 2.5 % was recorded in control followed by treatment two with 1.00 % none was recorded in the other treatment ($P < 0.05$). Results on the breast muscle revealed that saturated and unsaturated fatty acid was 29.80 %, 41.83 % in control, treatment two (28.11 %, 55.94 %), three (27.19 %, 56.97 %), four (23.15 %, 73.69 %) and five (22.90 %, 73.74 %) in each case ($P < 0.05$). Immunoglobulin A, G and M values were influenced ($P < 0.05$) by the treatments. In conclusion, feeding *Muntingia calabura* leaf extract up to 1.2 mL/liter of water had no negative effect on the performance and health status of birds.

*Corresponding author

Alagbe Olujimi John, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Received: July 08, 2024; **Accepted:** July 12, 2024; **Published:** July 22, 2024

Keywords: *Muntingia Calabura* Leaf Extract, Growth, Immune Response, Phytochemicals, Medicinal Plants, Food Safety

Introduction

The topic of antibiotic growth promoters and their replacement continues to be of great interest to poultry producers with a wide range of options now available to those who are either prevented from using antibiotic growth promoters by local regulations, or prefer to use an alternative due to their non-toxic effect. The use of medicinal plants is among the key alternative solutions to antibiotics because they possess phytochemicals that has high therapeutic properties and no withdrawal period [1,2].

Muntingia calabura popularly known as 'Jamaican cherry' is a rapidly growing small, evergreen plant belonging to family muntingiaceae and order – Malvales. The plant is native to southern Mexico central America and western south America. It is also widely cultivated in Asia including India [3]. The leaves are soft evergreen and alternate, ablong phyllotaxy about 4-15 cm long and 1-6 cm wide, oblong with toothed margin oblique at the base and characterized with dark green color and upper surface indicates presence of minute hair [4].

Reports on the pharmacological evaluation of *Muntingia calabura* leaf has shown that it possesses numerous phyto-constituents such as: flavonoids, terpenoids, tannins, alkaloids, saponins, amongst others which performs multiple biological activities:

antimicrobial, antioxidant, anti-helminthic, antifungal, antiviral, hepato-protective, antibacterial, anti-tumor, cytotoxic, anti-diuretic and hypolipidemic [5,6]. *Muntingia calabura* is popular in folklore medicine and has been used for the treatment of severe diarrhea, dysentery, urethral discharge, gastro-intestinal infection, headache, dermatitis and other bacterial infections [7,8].

The aqueous extract of leaf has been reported to contain volatile compounds such as myrcene, thymol, α terpinol, linalool, geraniol, nerol, citronellol, eugenol, α lonone, β sitosterol, α Amyrin, Lupelol, α tocopherol and β carotene [9,10]. In a study performed by the ethanolic extract of the different parts of *Muntingia calabura* demonstrated antimicrobial activity against *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans*, *Streptococcus* spp, *Salmonella* spp and *Aspergillus* spp [11]. It also showed that the extracts from their flowers demonstrated more antiseptic and antiplasmodic properties than the other parts of the plant.

Previous findings by notable researchers have shown that dietary supplementation of medicinal plants in broilers can positively influence their growth, gastro intestinal tract morphology and physiology immune response and meat quality [12-16]. However, there is little or no report on the dietary supplementation of *Muntingia calabura* leaf extract in broilers. This information is timely, because livestock producers have turned against the use of antibiotics due to the increasing cases of antimicrobial resistance and their health effect. These studies will further promote organic

agriculture and food safety.

Therefore, this study was carried out to ascertain the effect of feeding Muntingia calabura leaf extract on the growth performance, carcass characteristics, immune response and meat quality of broilers.

Materials and Methods

Experimental Station, Ethical Approval and Extraction of Muntingia Calabura

The trial was conducted at Sumitra Institute's poultry unit situated between 26° 17' N and 70° 47' E. Experiment was executed in accordance to the specifications laid down by ethics council of Sumitra Research Institute, Gujarat India in the month of September to November, 2023 (CN/117H/2023).

Fresh leaves of Muntingia calabura were collected within Sumitra Institute, Gujarat in the month of September, 2023 and sent to the Crop Science section for proper identification by a certified taxonomist (Dr. Singh Liu) for proper authentication. It was later assigned a reference number (JP/008/223), shade dried for six days with an electronic blender. Three hundred grams of grounded Muntingia calabura was soaked in 1 liter and boiled at seventy °C for ten minutes, kept for three days before it was sieved using a Whatman filter paper to obtain Muntingia calabura leaf extract and kept in the refrigerator at four °C. 200 milliliters of Muntingia calabura leaf extract was sent to the department of biochemistry, Sumitra Research Institute, Gujarat for further evaluation.

Animal Management, Experimental Diet and Design

500 1-day old broiler chicks (Cobb 500) were purchased from a commercial farm in Gujarat in the month of September, 2023 and moved to the poultry unit of Sumitra Research Institute, India. On arrival, birds were unboxed and the average initial weight was recorded using a digital scale before it was randomly distributed into five treatments of five replicates consisting of ten birds each. Birds were raised in an environmentally controlled battery cage equipped with automatic nipple drinker's, feeders measuring 200 cm by 150 cm by 60 cm (length × width × height) in three tiers per row and given anti-stress (Glucomol wsp® at 10 grams/10 liter). Continuous lighting system was provided to enable birds to have unrestricted access to feed and water. Basal diet was sufficient in all nutrients according to the recommendation of National Research Council in 1994. Birds in treatment one was fed basal diet without Muntingia calabura leaf extract while treatment two, three, four and five were fed same diet with Muntingia calabura leaf extract at 10 milliliters, 20 milliliters, 30 milliliters and 40 milliliters per liter of water correspondingly. A completely randomized design was adopted throughout the trial which lasted for sixty days.

Collected Data

Body weight gain was calculated as the difference between the final body weight and initial body weight. Feed consumption was determined as the difference between the feed offered and refused. Feed conversion ratio (feed consumed to produce a unit of gain) was computed as the ratio of average daily feed intake to average daily weight gain. Average daily feed intake was estimated by dividing the total feed intake by the duration of experiment in days while average daily weight gain was computed by dividing the weight gain by the duration of experiment in days, all the variables were expressed in grams.

Estimation of Phyto-Constituents in Muntingia Calabura Leaf Extract

The laboratory procedures for the determination of tannins, flavonoids, terpenoids, alkaloids, saponins and phenolic compounds were carried out according to methods outlined by Alagbe.

Proximate Analysis of Starter and Finisher Diet

Experimental diets were analyzed using Foss NIRSTM instrument which uses optical near infra-red technology to ensure precision in result. One hundred and fifty grams of feed sample was placed in the collection pan and the machine was adjusted at a wavelength precision of 400 – 2500 nm, optical band width (9.00 ± 0.2 nm), spectral resolution (0.5 nm) to give a faster result at an analysis time less than one minutes.

Carcass Evaluation and Measurement

At the end of the trial, ten birds were randomly selected from each treatment for carcass evaluation. Birds were feed starved for 12 hours, placed on clean water and slaughtered manually with a sharp knife and de-feathered by immersing each bird in hot water boiled at eighty °C for fifteen minutes. Carcass were eviscerated removing the internal organs and measured with using a digital scale. Back, wing, thigh, breast and drumsticks were measured with a digital scale as part of the carcass measurement. The dressing percentage was calculated as the proportion of carcass weight to slaughter weight multiplied by hundred.

Fatty Acid Analysis

Meat from the breast muscle of slaughtered chicken were collected into a sample bottle and placed in an ice pack before it was transferred to the laboratory for analysis. Fatty acid compositions were determined using a commercial kit (Fin analyzer, model – YLB 500 GC, China) following the specified procedures outlined by the manufacturer. 0.5 sodium hydroxide plus 1.5 milliliter methanol was added to each sample and mixed together followed by heating at 100°C for 5 minutes before it was allowed to cool down at 30 – 40 °C. Addition of trifluoroborane methanol solution allows the sample to form separate layers before standardization using YLB 500 GC which has the following technical specifications: Injection volume (1 µL), oven and capillary temperature (140°C, 250°C) and detector FID (100°C).

Immunoglobulin Examination

2 mL of blood samples were collected from ten birds were randomly selected from each treatment at the end of the experiment for immunoglobulin assessment. Test was carried out using a commercial diagnostic kit Immulite® 1000 system. To ensure precision machine was adjusted according to the manufacturer's recommendation, operating temperature was set at 18-32 °C while wavelength band (1000 nm).

Data Analysis

Data collected on growth performance, carcass evaluation and fatty acid analysis were analyzed using the general linear model procedures of Statistical Analysis Systems software (SAS). Differences between treatment means were separated using software of the same package. Significant differences were declared at $P < 0.05$

Experimental Outcome

The presence of tannins (502.58 milligram/gram), flavonoids (820.69 milligram/gram), terpenoids (401.17 milligram/gram), phenolic compounds (1006.5 milligram/gram), alkaloids (115.16

milligram/gram) and saponins (100.8 milligram/gram) were identified in Muntingia calabura leaf extract (Table 1). Phenolic compounds had the greater concentration while alkaloids had the least concentration in order of abundance.

As revealed in Table 2, final body of birds at 0-28 days (starter phase) took the form of 912.5 grams to 1084.9 grams, body weight gain (866.4 – 1084.9 grams per bird), average daily weight gain (30.94 to 38.75 grams per bird), total feed intake (1300.4 – 1300.9 grams), average daily feed intake (46.44 – 46.46 grams per bird), feed conversion ratio (1.20 – 1.50) and mortality (1.00 – 1.50 %). Body weight gain of birds fed diet 2 (0.3 mL Muntingia calabura leaf extract) and diet 3 (0.6 mL Muntingia calabura leaf extract) were similar ($P>0.05$) to those given diet 4 (0.9 mL Muntingia calabura leaf extract) and diet 5 (1.2 mL Muntingia calabura leaf extract) but significantly greater than those fed diet 1 (without Muntingia calabura leaf extract). Muntingia calabura leaf extract had no significant ($P>0.05$) on average daily feed intake while highest ($P<0.05$) mortality was recorded in birds given diet 1 (1.50 %) followed by diet 2 (1.0 %) while none was recorded among birds in diet 3, 4 and 5 correspondingly which also had an excellent feed conversion ratio. At the finisher phase (29 -56 days), average daily weight gain of birds fed diet 4 (0.9 mL Muntingia calabura leaf extract) were similar ($P>0.05$) to those in diet 5 (1.2 mL Muntingia calabura leaf extract) while those given diet 2 (0.3 mL Muntingia calabura leaf extract) and diet 3 (0.6 mL Muntingia calabura leaf extract) also followed similar trend ($P>0.05$) but significantly higher than those fed diet 1 (without Muntingia calabura leaf extract). Conversely, average daily feed intake was greater among birds fed diet 3, 4 and 5, intermediate in diet 2 and lowest in diet 1 ($P<0.05$). In the overall production cycle (0 – 56 days), average daily weight gain values which varied from 34.80 – 47.25 grams per bird were higher among birds fed diet 4 and 5, intermediate in diet 2 and 3, lowest in diet 1 ($P<0.05$). Average daily feed intake was greater among birds fed Muntingia calabura leaf extract relative to those fed diet 1 ($P<0.05$) which also has the highest mortality of 2.5 % at the end of the production cycle. Feed conversion ratio was significantly ($P<0.05$) impacted by feeding Muntingia calabura leaf extract to birds.

Effect of Muntingia calabura leaf extract on the carcass characteristics of broiler chicken (Table 4) revealed that slaughter weight, carcass weight and dressing percentage which took the form of 1971.2 – 2806.3 grams, 1421.5 to 2306.5 grams and 72.11 – 82.19 percent were higher among birds fed diet 4 (0.9 mL Muntingia calabura leaf extract) and diet 5 (1.2 mL Muntingia calabura leaf extract), intermediate among birds given diet 3 (0.3 mL Muntingia calabura leaf extract) and diet 2 (0.3 mL Muntingia calabura leaf extract), lowest in diet 1 (without Muntingia calabura leaf extract) ($P<0.05$). Average weights of heads, wing, liver and heart were exactly comparable ($P>0.05$) in all the groups except for those of legs, breast, thigh, back and gizzard ($P<0.05$).

Effect of Muntingia calabura leaf extract on the fatty acid characterization of meat in the breast muscle of broilers (Table 5) revealed at it contained saturated fatty acid which comprises of myristic acid, palmitic acid, stearic acid, arachidic acid, behenic acid and lignoceric acid values which vary from 2.50 – 3.81 %,

2.31 – 3.00 %, 10.02 – 13.81 %, 5.06 – 5.55 %, 1.82 – 2.71 % and 0.40 – 0.92 % in that order. Palmitoleic acid, oleic acid, erucic acid, gadoleic acid and nervonic acid (monounsaturated fatty acid) took the form of (2.16 – 5.17 %), (2.25 – 3.80 %), (10.64 – 12.71 %), (4.26 – 7.18 %) and (1.55 – 2.87 %) accordingly and polyunsaturated fatty acid; alpha (α)-linolenic acid (10.22 – 20.11 %), gamma (γ) linolenic acid (0.87 – 2.88 %), arachidonic acid (6.11 – 10.56 %), eicosadienoic acid (0.55 – 1.95 %), docosapentaenoic acid (1.97 – 3.84 %) and decosatraenoic acid (1.25 – 2.67 %). Muntingia calabura leaf extract had a significant ($P<0.05$) on all the variables examined in the experiment.

Effect of Muntingia calabura leaf extract on the Immunoglobulins of broilers presented in Table 6 showed that immunoglobulin A (Ig A), IgG and IgM took the form of 7.66 – 10.26 ($\mu\text{g/mL}$), 2.29 – 4.15 ($\mu\text{g/mL}$) and 1.29 – 2.98 ($\mu\text{g/mL}$) in that order. Ig A values were greater among birds fed diet 4 (0.9 mL Muntingia calabura leaf extract) and diet 5 (1.5 mL Muntingia calabura leaf extract/litre of water), intermediate in diet 2 (0.3 mL Muntingia calabura leaf extract per litre of water) and diet 3 (0.6 mL Muntingia calabura leaf extract) and lowest in diet 1 (without Muntingia calabura leaf extract) ($P<0.05$). Conversely, IgG and IgM values were higher among birds fed Muntingia calabura leaf extract relative to those without Muntingia calabura leaf extract in diet 1 ($P<0.05$).

Table 1:Ingredients and Chemical Composition of Experimental Diet

	Starter phase (0 – 28 d)	Finisher phase (29 – 56 d)
Ingredients	Quantity (%)	Quantity (%)
Yellow corn	53.00	55.95
Wheat bran	5.00	6.00
Soybean meal	26.50	25.00
Groundnut meal	5.00	4.00
Fish meal (Imported: 72%)	5.00	2.00
Limestone	1.50	2.00
Bone meal	3.00	4.00
Methionine	0.25	0.25
Lysine	0.20	0.25
Mineral/Vitamin Premix	0.25	0.25
Salt	0.25	0.25
Toxin binder	0.05	0.05
Total	100.0	100.0
Analyzed analysis (% dry matter)		
Crude protein (%)	23.47	21.16
Crude fibre (%)	3.56	4.82
Ether extract (%)	3.20	3.83
Calcium (%)	1.49	1.47
Phosphorus (%)	0.58	0.57
Metabolizable energy (MJ/kg)	11.89	12.30

Table 2: Phyto-Constituents in Muntingia Calabura Leaf Extract

Chemical components	Units	Concentration
Tannins	Milligram/gram	502.58
Flavonoids	Milligram/gram	820.69
Terpenoids	Milligram/gram	401.17
Phenols	Milligram/gram	1006.5
Alkaloids	Milligram/gram	115.16
Saponins	Milligram/gram	200.8

Table 3: Effect of Muntingia Calabura Leaf Extract on the Growth Performance of Broiler Chicken

0-28 day	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P-value
Number of birds	60.00	60.00	60.00	60.00	60.00	-	
Initial body weight (grams/bird)	46.10	45.80	45.60	45.50	46.00	0.51	0.26
final body weight (grams/bird)	912.5 ^b	1120.7 ^a	1120.9 ^a	1130.1 ^a	1130.9 ^a	29.81	0.38
Body weight gain (grams/bird)	866.4 ^b	1074.9 ^a	1075.3 ^a	1084.6 ^a	1084.9 ^a	27.26	0.61
Average daily weight gain (grams/bird)	30.94 ^b	38.39 ^a	38.40 ^a	38.74 ^a	38.75 ^a	0.41	0.19
Total feed intake (grams/bird)	1300.3	1300.5	1300.6	1300.8	1300.9	31.83	0.37
Average daily feed intake (grams/bird)	46.44	46.45	46.45	46.45	46.46	0.50	0.25
Feed conversion ratio	1.50 ^a	1.21 ^b	1.20 ^b	1.20 ^b	1.20 ^b	0.07	0.03
Mortality (percentage)	1.50 ^a	1.00 ^b	0	0	0	0.02	0.01
29-56 days							
Body weight gain (grams/bird)	1902.7 ^c	2300.8 ^b	2319.5 ^b	2509.8 ^a	2600.1 ^a	51.42	0.02
Average daily weight gain (grams/bird)	67.95 ^c	82.17 ^b	82.83 ^b	90.64 ^a	92.86 ^a	0.72	0.06
Total feed intake (grams/bird)	3410.8 ^c	3610.2 ^b	3650.8 ^a	3655.1 ^a	3658.1 ^a	61.21	0.08
Average daily feed intake (grams/bird)	121.8 ^c	128.9 ^b	130.4 ^a	130.5 ^a	130.6 ^a	0.62	0.01
feed conversion ratio	1.80 ^a	1.57 ^b	1.57 ^b	1.46 ^c	1.41 ^d	0.06	0.01
Mortality (percentage)	1.00	0	0	0	0	0.02	0.01
Overall production cycle (0-56 days)							
Body weight gain (grams/bird)	1948.8 ^c	2346.6 ^b	2365.1 ^b	2555.3 ^a	2646.1 ^a	49.82	0.06
Average daily weight gain (grams/bird)	34.80 ^c	42.00 ^b	42.23 ^b	46.00 ^a	47.25 ^a	0.42	0.30
Total feed intake (grams/bird)	4711.1 ^b	4910.7 ^a	4951.4 ^a	4955.9 ^a	4959 ^a	105.2	0.03
Average daily feed intake (grams/bird)	84.13 ^b	87.69 ^a	88.42 ^a	88.50 ^a	88.55 ^a	0.70	0.06
Feed conversion ratio	2.42 ^a	2.09 ^b	2.09 ^b	1.94 ^c	1.87 ^d	0.03	0.04
Total mortality (percentage)	2.50 ^a	1.00 ^b	0	0	0		

Table 4: Effect of Muntingia Calabura Leaf Extract on the Carcass Characteristics of Broilers

Variables	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P-value
Slaughter weight (grams)	1971.2 ^c	2435.7 ^b	2441.5 ^b	2804.1 ^a	2806.3 ^a	28.22	0.42
Carcass weight (grams)	1421.5 ^c	1835.6 ^b	1821.9 ^b	2304.3 ^a	2306.5 ^a	23.97	0.30
Dressing percentage	72.11 ^c	75.36 ^b	74.62 ^b	82.18 ^a	82.19 ^a	0.90	0.02
Head (grams)	8.02	8.06	8.40	8.38	8.51	0.05	0.01
Legs (grams)	9.76 ^c	10.51 ^b	10.60 ^b	12.22 ^a	12.96 ^a	0.07	0.02
Breast (grams)	411.7 ^c	563.1 ^b	591.0 ^b	709.8 ^a	715.1 ^a	9.11	0.18
Thigh (grams)	310.0 ^b	314.1 ^b	306.3 ^b	490.1 ^a	495.7 ^a	4.38	0.21
Back (grams)	183.2 ^c	203.6 ^b	200.0 ^b	259.2 ^a	261.4 ^a	4.01	0.08
Wing (grams)	79.0	78.8	79.5	80.8	82.1	0.72	0.01
Drumstick (grams)	51.3 ^c	63.5 ^b	65.2 ^b	71.6 ^a	72.9 ^a	0.65	0.02
Liver (grams)	31.2	33.5	32.8	33.1	33.0	0.45	0.04
Gizzard (grams)	30.6 ^c	38.7 ^b	39.5 ^b	43.9 ^a	45.1 ^a	0.08	0.03
Heart (grams)	12.6	12.9	13.2	13.8	13.1	0.02	0.04

Means on the same row having different superscripts are significantly different (P<0.05); SEM: Standard Error of Mean; Diet 1: basal diet without Muntinda calabura; diet 2: basal diet with 0.3 mL Muntinda calabura leaf extract/litre of water; diet 3: basal diet with 0.6 mL Muntinda calabura leaf extract/litre of water; diet 4: basal diet with 0.9 mL Muntinda calabura leaf extract/litre of water; diet 5: basal diet with 1.2 mL Muntinda calabura leaf extract/litre of water

Table 5: Effect of Muntingia Calabura Leaf Extract on the Fatty Acid of Breast Muscle of Broilers

Variables	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P-value
Saturated fatty acids							
Myristic acid	3.81 ^a	2.90 ^b	2.61 ^b	2.56 ^b	2.50 ^b	0.82	0.04
Palmitic acid	3.00 ^a	2.71 ^b	2.62 ^b	2.31 ^b	2.26 ^b	0.19	0.33
Stearic acid	13.81 ^a	13.74 ^b	13.83 ^b	10.91 ^c	10.02 ^c	0.21	0.10
Arachidic acid	5.55	5.45	5.22	5.06	5.9	0.23	0.11
Behenic acid	2.71 ^a	2.60 ^b	2.22 ^b	1.87 ^c	1.82 ^c	0.30	0.40
Lignoceric acid	0.92 ^a	0.71 ^b	0.69 ^b	0.44 ^c	0.40 ^c	0.18	0.50
Monounsaturated fatty acid							
Palmitoleic acid	2.16 ^c	3.88 ^b	3.73 ^b	5.11 ^a	5.17 ^a	0.66	0.04
Oleic acid	2.25 ^c	3.00 ^b	3.03 ^b	3.72 ^a	3.80 ^a	0.15	0.06
Erucic acid	10.64 ^b	10.22 ^b	10.70 ^b	12.02 ^a	12.71 ^a	0.35	0.01
Gadoleic acid	4.26 ^c	5.00 ^b	5.06 ^b	7.04 ^b	7.18 ^a	0.27	0.12
Nervonic acid	1.55 ^c	2.20 ^b	2.33 ^b	2.80 ^a	2.87 ^a	0.01	0.05
Polyunsaturated fatty acids							
Alpha-linolenic acid	10.22 ^c	15.00 ^b	15.27 ^b	20.09 ^a	20.11 ^a	0.09	0.40
Gamma linolenic acid	0.87 ^c	1.90 ^b	1.92 ^b	2.80 ^a	2.88 ^a	0.52	0.00
Arachidonic acid	6.11 ^c	9.30 ^b	9.38 ^b	12.41 ^a	10.56 ^a	0.06	0.01
Eicosadienoic acid	0.55 ^c	0.90 ^b	0.93 ^b	1.91 ^a	1.95 ^a	0.28	0.00
Docosapentaenoic acid	1.97 ^c	2.46 ^b	2.57 ^b	3.21 ^a	3.84 ^a	0.23	0.10
Decosatetraenoic acid	1.25 ^c	2.00 ^b	2.05 ^b	2.58 ^a	2.67 ^a	0.13	0.00
Total saturated fatty acid (%)							
Total saturated fatty acid (%)	29.8 ^a	28.11 ^a	27.19 ^b	23.15 ^c	22.9 ^c	0.96	0.04
Total unsaturated fatty acid (%)							
Total unsaturated fatty acid (%)	41.83 ^c	55.94 ^b	56.97 ^b	73.69 ^a	73.74 ^a	1.83	0.02

Means on the same row having different superscripts are significantly different (P<0.05); SEM: Standard Error of Mean; Diet 1: basal diet without Muntinda calabura; diet 2: basal diet with 0.3 mL Muntinda calabura leaf extract/litre of water; diet 3: basal diet with 0.6 mL Muntinda calabura leaf extract/litre of water; diet 4: basal diet with 0.9 mL Muntinda calabura leaf extract/litre of water; diet 5: basal diet with 1.2 mL Muntinda calabura leaf extract/litre of water

Table 6: Effect of Muntingia Calabura Leaf Extract on the Immunoglobulins of Broilers

Variables	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	P-value
Immunoglobulin A (µg/mL)	7.66 ^c	9.06 ^b	9.33 ^b	10.10 ^a	10.26 ^a	0.96	0.51
Immunoglobulin G (µg/mL)	2.29 ^b	3.31 ^a	3.38 ^a	4.05 ^a	4.15 ^a	0.05	0.04
Immunoglobulin M (µg/mL)	1.93 ^b	2.72 ^a	2.79 ^a	2.95 ^a	2.98 ^a	0.07	0.01

Means on the same row having different superscripts are significantly different (P<0.05); SEM: Standard Error of Mean; Diet 1: basal diet without Muntinda calabura; diet 2: basal diet with 0.3 mL Muntinda calabura leaf extract/litre of water; diet 3: basal diet with 0.6 mL Muntinda calabura leaf extract/litre of water; diet 4: basal diet with 0.9 mL Muntinda calabura leaf extract/litre of water; diet 5: basal diet with 1.2 mL Muntinda calabura leaf extract/litre of water

Discussion

In the overall production cycle, the higher average daily weight gain was highest among birds fed diet 4 (0.9 mL Muntingia calabura leaf extract per liter of water) and diet 5 (1.2 mL Muntingia calabura leaf extract per liter of water) can be attributed to the presence of phyto-constituents in Muntingia calabura leaf extract as presented in Table 2. According to this study, Muntingia calabura leaf extract contains phenolic compounds, terpenoids, flavonoids, alkaloids, saponins and tannins at different concentrations, these

compounds have been reported to possess numerous medicinal or herbal properties [17]. This result confirms the earlier study of who reveals that Muntingia calabura leaves contains flavonoids, alkaloids, tannins and phenols which have antimicrobial, anti-inflammatory, antioxidant, antifungal, antiviral, cytotoxic, anti-helminthic, analgesics, anti-diuretic, amongst others [18-21]. A synergistic combination of these phyto-constituents especially when administered at 0.9 mL (diet 4) and 1.2 mL (diet 5) per liter of water was capable of stimulating the secretion of digestive enzymes and saliva to improve the overall nutrient digestibility in birds [14,22]. The antimicrobial properties in Muntingia calabura leaves also explains the reason why mortality was not recorded among birds fed the diet 3, 4 and 5. The presence of phenols, which is the most abundant phyto-constituents in this study could strengthen the gut epithelial barrier, thereby making it less prone to pathogenic organisms [23]. Best feed conversion ratio was recorded among birds fed diet 3 and 4 relative to the other groups. The result recorded in this experiment is in consonance with the reports of when plant extracts derived from thyme was fed to broilers at 2 mL/litre of water [12]. recorded similar results when phytochemicals were fed to broiler chicken. Conversely, recorded a non-significant difference in the feed intake of broilers fed *Ocimum gratissimum* leaf powder, this discrepancy could be attributed to the levels supplemented in the diet as well as the nature of bioactive compound in the test ingredient [24-26].

The highest dressing percentage were found in birds fed 0.9 mL and 1.2 mL Muntingia calabura leaf extract per liter of water in diet 4 and 5, intermediate in 0.3 mL and 0.6 mL Muntingia calabura leaf extract per liter of water and lowest in diet 1 (without Muntingia calabura leaf extract) [25]. The current results are in agreement with the previous finding of who reported that dressing percentage increased when broilers were fed 2.5 g/kg *Ocimum gratissimum* leaf powder as compared to the control [25]. Similarly, reported that dressing percentages were significantly increased when 0.5, 1.0 and 1.5% turmeric powder was included in the diets of broiler chickens. In disagreement with this current experiment, supplemented *Nigella sativa* seeds and fresh garlic at 1.5g/kg and it revealed no significant changes in carcass weight and dressing percentage. Weights of head, wing, liver and heart were not significantly ($P>0.05$) influenced by the treatment except for those of legs, breast, thigh, back and drumstick ($P>0.05$), this result removes the possibility of toxicity from Muntingia calabura leaf extract since organs were not enlarged [27,28]. The increase in thigh, breast, gizzard and breast could be due to the presence of the phyto-constituents in the test ingredient [29]. Gizzard mechanically grinds food and acts as a microbial barrier due to its low pH and an increase suggests an improvement in digestive or metabolic capacity of birds [30].

Meat quality can be influenced by many factors such as management, genetics, environment and nutrition [31,32]. Total saturated fatty acid and total unsaturated fatty acid took the form of 29.80 – 22.90 % and 41.83 – 73.74 % accordingly was within the range reported by when natural antioxidant was supplemented in the diet of broiler chicken. Similar outcome was observed by when quercetin was supplemented in the diet of broilers [33-35]. Unsaturated fatty acid (monosaturated fatty acid and polyunsaturated fatty acid) decreases the risk of heart disease by reducing the amount of low-density lipoprotein (bad cholesterol) [LDL] and increasing high density lipoprotein [HDL]. Conversely, saturated fatty acid increases both LDL and HDL [15]. Result obtained revealed that birds fed Muntingia calabura leaf extract had higher concentrations of unsaturated fatty acid and low saturated fatty acid relative to the control suggesting that the test ingredient

has a tendency to lower cholesterol (hypolipidemic) in meat thus promoting food safety and increasing consumer's acceptability [20]. This is made possible by the presence of phytochemicals such as flavonoids and phenols in Muntingia calabura leaf extract. According to flavonoids and phenols are capable of reducing the risk of hyperlipidemia in the serum and meat of animals.

Results on the effect of Muntingia calabura leaf extract on the Immunoglobulins of broilers showed that it influenced the activities of immunoglobulin A (IgA), IgG and IgM in birds compared to those given control diet. The results suggest that Muntingia calabura leaf extract possess immune-modulatory properties thus reducing the exposure of birds to pathogenic attack. According to leaves of Muntingia calabura contains bioactive compounds like; 20,40- dihydroxy chalcone (isoliquiritigenin (cabrevin), (2S)-50-hydroxy-7,8,30,40-tetramethoxyflavan 20,40-dihydroxydihydrochalcone and 3,4,5-trihydroxybenzoic acid which are strong antioxidants capable of producing antibodies in the system of birds [19]. Immunoglobulin G has been reported to facilitate the process of phagocytosis and provides immunity to the system of animals while IgA and IgM are capable of protecting the mucosal membrane and blood stream to limit inflammation in the body [36]. Result obtained in this study is in agreement with the reports of who observed an increase in the immunoglobulins of broilers fed herb mixture consisting of *Zingiber officinale*, *Platycodon radix* and *Phlomis umbrosa* [37].

Conclusion

In conclusion, Muntingia calabura leaf extract have beneficial effects on birds because they possess phyto-constituents that has antimicrobial, anti-inflammatory, sensorial stimulation, flavoring, immuno-stimulatory properties amongst others. Supplementing Muntingia calabura leaf extract at 0.9 mL and 1.2 mL per litre of water reveals that it can optimize the performance of birds without compromising the meat quality as well as the health status of animals. This research will further help to reduce the increasing rate of multidrug resistance and promote food safety [38-48].

References

1. John AO, Muritala SD, Aduragbemi YA, Kadiri CM, Bamiloye SO, et al. (2024) The approximate mineral and phytochemical content of the leaves, stem bark and roots of *Pterocarpus erinaceus* in India Cerrado. *Agricultural and Biological Research* 1: 32-41.
2. Ojediran TK, Emiola IA, Durojaye V, John Alagbe (2024) Proximate, vitamin and GC-MS profiling of *Kigella africana* powder Cerrado. *Agricultural and Biological Research* 1: 13-20.
3. Sani MH, Zakaria ZA, Balan T, The LK, Salleh MZ (2012) Antinociceptive activity of methanol extract of Muntingia calabura leaves and the mechanisms of action involved. *Evidence Based Complementary Alternative Medicine* 2012: 890361.
4. Su BN, Park EJ, Vigo JS, Graham JG, Cabieses F, et al. (2003) Activity-guided isolation of the chemical constituents of Muntingia calabura using a quinine reductase induction assay. *Journal of Phytochemistry* 63: 335-341.
5. Chen JJ, Lee HH, Shih CD, Liao CH, Chen IS, et al. (2007) New dihydrochalcones and anti-platelet aggregation constituents from the leaves of Muntingia calabura. *Planta Medicine* 73: 572-577.
6. Chen JJ, Lin RW, Duh CY, Huang HY, Chen IS (2004) Flavones and cytotoxic constituents from the stem bark of Muntingia calabura. *Journal of Chinal Chemical Society* 51: 665-670.

7. Balan T, Mohd Sani MH, Suppaiah V, Mohtarrudin N, Suhaili Z, et al. (2013) Antiulcer activity of *Muntingia calabura* leaves involves the modulation of 1385 endogenous nitric oxide and nonprotein sulphhydryl compounds. *Pharmaceutical Biology* 52: 410-418.
8. Zakaria ZA, Sulaiman MR, Jais AMM, Somchit MZ, Jayaraman KV (2006a) The antinociceptive activity of *Muntingia calabura* aqueous extract and the involvement of L-arginine/nitric oxide/cyclic guanosine monophosphate pathway in its observed activity in mice. *Fundamental Clinical Pharmacology* 20: 365-372.
9. Preethi K, Vijayalakshmi N, Shamna R, Sasikumar JM (2010) In vitro antioxidant activity of extracts from fruits of *Muntingia calabura* Linn from India. *Pharmacognosy J* 2: 11-18.
10. Zakaria ZA, Fatimah CA, Jais AMM, Zaiton H, Henie EFP, et al. (2006b) The in vitro antibacterial activity of *Muntingia 1544 calabura* extract. *Journal of Pharmacology* 2: 290-293.
11. Ibrahim IAA, Abdulla MA, Abdelwahab SI, Al-Bayaty F, Majid NA (2012) Leaves extract of *Muntingia calabura* protects against gastric ulcer induced by ethanol in 1430 Sprague-dawley rats. *Clinical Experimental Pharmacology* S5: 004.
12. Alipour F, Hassanabadi A, Golian A, Nassiri-Moghaddam H (2015) Effect of plant extracts derived from thyme on male broiler performance. *Poultry Science* 94: 2630-2634.
13. Oloruntola OD, Agbede JO, Ayodele SO, Oloruntola DA (2018a) Neem, pawpaw, and bamboo leaf meal dietary supplementation in broiler chickens: Effect on performance and health status. *Journal of Food Biochemistry* e12723.
14. Alagbe Olujimi John (2024) Novel phytochemicals' impact on weaned pigs' growth performance, haematology and serum biochemical indicators. *Black Sea Journal of Agriculture* 7: 82-89.
15. Mgbemena MM, Ilechukwu I, Okwunodolu FU, Chukwurah JO, Lucky IB (2019) Chemical composition, proximate and phytochemical analysis of *Irvingia gabonensis* and *Irvingia qombolu* peels, seed coat, leaves, and seeds. *Ovidius University Annals of Chemistry* 30: 65-69.
16. Gou FC, Williams BA, Kwakkel RP, Li ZP, Lou JY, et al. (2004) Effects of mushroom and herb polysaccharides as alternatives for an antibiotic, on the cecal microbial ecosystem in broiler chickens. *Poultry Science* 83: 175-182.
17. Adewale AO, Alagbe JO, Adeoye Adekemi O (2021) Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytochemical Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies* 3: 1-12.
18. IMahmood ND, Nasir NL, Rofiee M, Tohid S, Ching SM, et al. (2014) *Muntingia calabura*: A review of its traditional uses, chemical properties, and pharmacological observations. *Journal of Pharmaceutical and Biological Sciences* 52: 1598-1623.
19. Mohini U, Mohini K, Rohini P, Shailja K, Priya G (2021) *Muntingia calabura*: A comprehensive review. *Journal of Pharmaceutical and Biological Sciences* 9: 81-87.
20. Shittu MD, Alagbe JO, Adejumo DO, Ademola SG, Abiola AO, et al. (2021) Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. *Bioinformatics and Proteomics Open Access Journal* 5: 000143.
21. Agubosi OCP, Soliu MB, Alagbe JO (2022) Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. *Central Asian Journal of Theoretical and Applied Sciences* 3: 30-39.
22. John AO (2024) Growth performance, haemato-biochemical indices of broiler chicken fed *Aristochia indica* as a phytochemical feed additive Cerrado. *Agricultural and Biological Research* 1: 42-53.
23. Liz M (2020) The value of direct immune support as an alternative to antibiotic growth promoters. *International Poultry Magazine* 4: 5-7.
24. Olujimi JA (2024) Impact of *Cordyline fruticosa* leaf meal supplemented diet on growth performance, egg production, egg quality and some haematological indices of laying hen. *Discovery Agriculture* 10: e14da1570.
25. Oloruntola OD, Adu AO, Gbore FA, Falowo AB, Olarotimi OJ (2021) Performance of broiler chicken fed diets supplemented with *Irvingia gabonensis* and *Ocimum gratissimum* leaf powder. *Slovak Journal of Animal Science* 54: 7-20.
26. John AO (2024) *Clerodendron splendens* leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population Cerrado. *Agricultural and Biological Research* 1: 21-31.
27. El-Ghamry AA, El-Mallah GM, El-Yamny AT (2002a) The effect of incorporation yeast culture, *Nigella sativa* seeds and fresh garlic in broiler diets on their performance. *Egyptian Poultry Science* 22: 445-459.
28. Durrani FR, Ismail M, Sultan A, Suhail SM, Chand N, et al. (2006) Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agricultural and Biological Science* 1: 9-11.
29. Duru M, Zeynep E, Asuman D (2013) Effect of seed powder of a herbal legume Fenugreek (*Trigonella foenum-graceum* L.) on growth performance, body components, digestive parts, and blood parameters of broiler chicks. *Pakistan Journal of Zoology* 45: 1007-1014.
30. Mushtaq MH, Parvin R, Kim J (2014) Carcass and body organ characteristics of broilers supplemented with dietary sodium and sodium salts under a phase feeding system. *Journal of Animal Science and Technology* 56: 1-7.
31. Diana S (2021) Influence of amino acids arginine and histidine on broiler meat quality. *International Poultry Magazine* 2: 3-4.
32. Alagbe JO (2022) Use of medicinal plants as a panacea to poultry production and food security: A review. *Gospodarka I Innowacje* 22: 1-12.
33. Kumar Y, Yadav DN, Ahmad T, Narsaiah K (2015) Recent trends in the use of natural antioxidants for meat and meat products. *Comprehensive Review of Food Sciences and Safety* 14: 796-812.
34. Goliomytis M, Tsourekis D, Simitzis PE, Charismiadou MA, Hager T, et al. (2014) The effect of quercetin dietary supplementation on broiler growth performance, meat quality and oxidative stability. *Poultry Science* 93: 1957-1962.
35. Agubosi OCP, Alexander James, Alagbe JO (2022) Influence of dietary inclusion of Sunflower (*Helianthus annuus*) oil on growth performance and oxidative status of broiler chicks. *Central Asian Journal of Medical and Natural Sciences* 2: 187-195.
36. Begum M, Hosain MM, Kim IH (2014) Effect of plant extract on the growth performance meat quality, relative organ weights, nutrient digestibility and blood profiles in broiler chickens: possible role of insulin-like growth promoter. *Veterinarian Medicine* 59: 415-423.
37. Preethi K, Vijayalakshmi N, Shamna R, Sasikumar JM (2012) In vitro antioxidant activity of extracts from fruits of

- M calabura linn from India. *Pharmacognosy Journal* 2: 11-18.
38. Zakaria ZA, Mohamed AM, Jamil M, Rofiee NS, Hussain MS, et al. (2011) Vitro Antiproliferative and Antioxidant Activities of the Extracts of M. calabura Leaves. *American Journal of Chemical Medicine* 39: 183-200.
 39. Zakaria A, Sufian AS, Ramasamy K, Ahmat N, Sulaiman MR, et al. (2010) In vitro antimicrobial activity of Muntingia calabura extracts and fractions. *African Journal of Microbiology Research* 4: 304-308.
 40. National Research Council (1994) Nutrient Requirements of Poultry eighth ed. National Academic Press Washington DC.
 41. Alagbe John Olujimi, Ramalan Sadiq Muhammad, Shittu Muritala Daniel, Olagoke Olayemi Christiana (2022) Effect of Trichilia monadelpha stem bark extract on the fatty acid composition of rabbit's thigh meat. *Journal of Environmental Issues and Climate Change* 1: 63-71.
 42. Alagbe JO, Shittu MD, Tanimomo Babatunde K (2022) Influence of Anogeissusleio carpus stem bark on the fatty acid composition in meat of broiler chickens. *European Journal of Life Safety and Stability* 14: 13-22.
 43. Singh Sharma, Alagbe Olujimi John, Liu Xing, Sharma Ram, Kumar Amita (2022) Comparative analysis of ethanolic Juniperus thurifera leaf, stem bark and root extract using gas chromatography and mass spectrometry. *International Journal of Agriculture and Animal Production* 2: 18-27.
 44. John AO (2024) Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken. *Brazilian Journal of Science* 3: 82-95.
 45. Ojediran TK, Emiola IA, Durojaye V, John Alagbe (2024) Analysis of Kigellia africana fruit's powder antioxidant and phytochemical properties. *Brazilian Journal of Science* 3: 38-49.
 46. John AO (2024) Impact of dietary supplementation of Rhamnus prinoides leaf extract on the growth performance, nutrient retention and intestinal microbial count of Japanese quails. *Brazilian Journal of Science* 3: 40-50.
 47. John AO (2024) Effect of performance, serum biochemistry and heamatological components of feeding Japanese quails phytogetic feed additions comparising Megaphrynum macrostachyum leaves. *Brazilian Journal of Science* 3: 51-64.
 48. Elagib HAA, Abbas SA, Elamin KM (2013) Effect of different natural feed additives compared to antibiotic on performance of broiler chicks under high temperature. *Bulletin of Environmental and Pharmacological Life Science* 2: 139-144.

Copyright: ©2024 Alagbe Olujimi John. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.